



On increasing global temperatures: 75 years after Callendar

Ed Hawkins^{a*} and Phil. D. Jones^{b,c}

^aNCAS-Climate, Department of Meteorology, University of Reading, UK.

^bClimatic Research Unit, University of East Anglia, Norwich, UK.

^cCenter of Excellence for Climate Change Research, Department of Meteorology, King Abdulaziz University, Jeddah, Saudi Arabia.

*Correspondence to: Department of Meteorology, University of Reading, Reading. RG6 6BB. UK.
E-mail: e.hawkins@reading.ac.uk

In 1938, Guy Stewart Callendar was the first to demonstrate that the Earth's land surface was warming. Callendar also suggested that the production of carbon dioxide by the combustion of fossil fuels was responsible for much of this modern change in climate. This short note marks the 75th anniversary of Callendar's landmark study and demonstrates that his global land temperature estimates agree remarkably well with more recent analyses.

This is a preprint of an article accepted for publication in *Quarterly Journal of the Royal Meteorological Society* Copyright © 2013 Royal Meteorological Society

Key Words: global temperatures; history; Guy Stewart Callendar

Received 18 January 2013; Revised 12 April 2013; Accepted 15 April 2013

Citation: Hawkins E, Jones PD. 2013. On increasing global temperatures: 75 years after Callendar. *Q. J. R. Meteorol. Soc.* in press

1. Introduction

The discussion of the human influence on global temperatures has a long history. For instance, Fourier (1824) first discussed why the Earth was warmer than expected from solely considering solar radiation reaching the planet. Fourier ruled out geothermal effects, considered the temperature of outer space, and pondered about heat being trapped in the atmosphere, but did not come to any firm conclusions. Tyndall (1861) suggested a solution to this conundrum by experimentally demonstrating that gases such as carbon dioxide can effectively absorb infrared radiation, i.e. the 'greenhouse effect'.

Later, Arrhenius (1896) used the results of Tyndall and others to produce the first estimate of the sensitivity of global temperatures to increases in carbon dioxide. Much of this research was aimed at understanding historical climatic variations such as ice ages rather than any modern changes (e.g. Ekholm 1901). However, in the early 1900s, the theory that increases in atmospheric carbon dioxide could change the climate was not widely accepted.

Separately, widespread temperature records were being used to estimate the variability in global land air temperatures during the late 1800s (e.g. Köppen 1881), but

no trend was evident. These activities eventually culminated in the work of Callendar (1938) who was the first to show that the Earth's land temperature had increased over the previous 50 years. Callendar also suggested that the production of carbon dioxide by the combustion of fossil fuels was responsible for a large part of this warming, which became known for a time as the 'Callendar Effect' (see Fleming (2007) and Weart (2008) for further discussion).

This short note marks the 75th anniversary of Callendar's landmark study, originally published in the *Quarterly Journal of the Royal Meteorological Society* in April 1938.

2. Comparing global temperature estimates

As highlighted by Le Treut *et al.* (2007), many studies have produced consistent estimates of the smoothed decadal variability in global temperatures in the instrumental period (since about 1850). Here we compare the inter-annual variability of three of these land-based records as consistently as possible.

Fig. 1 compares the latest CRUTEM4 (Jones *et al.* 2012) estimates for annual near-global land temperatures with that of Callendar (1938). The agreements in trends and variability are striking - the correlation between the




Figure 1. Comparing historical reconstructions of near-global land temperatures using CRUTEM4 (black, Jones *et al.* 2012) with Callendar (1938) (red) and Callendar (1961) (blue), using a reference period of 1880–1935. The CRUTEM4 estimates are for 60°S–60°N (to accord with Callendar’s series), with grey shading representing the 95% uncertainty.

two timeseries during their common period is $r = 0.81$. However, CRUTEM4 shows larger amplitude variability. Callendar had access to temperatures from just 147 land-based stations to reconstruct his global temperature timeseries, whereas CRUTEM4 includes more than 200 in 1880, increasing to more than 2000 by 1935. The differing amplitudes of variability is therefore perhaps surprising as fewer stations might be expected to result in a higher variance (e.g. Jones *et al.* 1997), but this feature remains unexplained.*

The analysis is restricted to 60°S–60°N because Callendar did not include any Antarctic or Arctic stations in his record. However, he did note that the omitted Arctic stations had the largest variations and that the northern latitudes were warming faster than other regions. More recently, Wood *et al.* (2010) demonstrated that the high northern latitudes underwent a striking temperature rise during this period.

In addition, Callendar produced timeseries for different regions, including Western Europe, New York State, as well as different latitudinal bands. Sadly he did not publish the values for these, except for decadal averages of the Northern temperate zone. Callendar suggests that, “*quite a moderate number of reliable temperature records may be used to give the period anomalies of very large areas*”, a fact which has been discussed in a number of more recent papers (see e.g. Madden *et al.* 1993; Jones *et al.* 1997). The latter provide a quantification of the number required to derive reliable large-scale averages in terms of the expected number of spatial degrees of freedom. The number depends on the time scale of interest, but as few as 50 well-spaced locations should provide reliable estimates at the annual timescale.

*The data in the original paper are tabulated as annual means, so we have assumed this to be correct. However, it is possible that Callendar temporally smoothed some of the data, resulting in a lower variance.

Callendar later expanded his estimates of global and regional land temperatures using a much larger number of station records and also published the data for zonal and seasonal averages (Callendar 1961). A comparison of the extended near-global (60°S–60°N) annual mean dataset is also shown in Fig. 1, which agrees even better with CRUTEM4 ($r = 0.92$), and has a similar amplitude of variability.

3. Reasons for the observed temperature increase

A large part of Callendar (1938) discusses the change in global temperatures that would have been expected given the observed increases in atmospheric carbon dioxide concentration. The calculations were somewhat hindered by the existing understanding of atmospheric radiative physics, and by the limited available observations of the infrared absorption spectrum and carbon dioxide concentrations. In addition, he considered the energy balance at the surface instead of the top of the atmosphere. Nevertheless, Callendar estimated that global temperatures should have warmed by around 0.03 K/decade due to the increasing levels of carbon dioxide. The global temperature estimates exhibited a larger trend of 0.06 K/decade, leading to his conclusion that the combustion of fossil fuels was making a significant contribution to the observed warming. Perhaps Callendar was slightly fortunate that the observed rise in temperatures was larger than expected from purely anthropogenic effects, therefore making it easier to detect.

Callendar also considered other reasons for the temperature increase, notably the urban heat island (UHI) effect. He separated his sample into stations within cities and those in rural areas, and found “*no secular increase of temperature due to city influence*”, although his sample sizes were small. More recent estimates also suggest that the effect of the

UHI and exposure of thermometers on global temperatures is small (e.g. Parker 2010; Jones and Wigley 2010).

4. Discussion

Callendar's conclusions were treated with a degree of skepticism. He even starts his introduction by saying that, "[f]ew of those familiar with the natural heat exchanges of the atmosphere, which go into the making of our climates and weather, would be prepared to admit that the activities of man could have any influence upon phenomena of so vast a scale". The peer-review discussion section which appears with Callendar (1938) expresses concern about the radiative physics calculations and the accuracy of the carbon dioxide measurements used. In fact, Callendar's findings on the role of carbon dioxide were largely dismissed by the scientific community until improved estimates of the infrared absorption spectrum were obtained and analysed (e.g. Callendar 1941; Plass 1956).

There was also debate about whether the observed temperature increase was in fact a natural fluctuation or caused by changes in atmospheric circulation, rather than an anthropogenic fingerprint. Indeed, there is still much discussion (e.g. Delworth and Knutson 2000; Bronnimann 2009) over the causes of this early twentieth century warming, which may have had considerable spatial and seasonal variations. However, it seems likely that there were multiple causes, including an internal climate fluctuation, recovery from the large volcanic forcing at the end of the 19th century, as well as solar and anthropogenic influences.

We also note that additional weather observations are even now becoming available from many previously non-digitized historical records, including the period covered by Callendar. For example, the Old Weather project (<http://www.oldweather.org>) has recently completed the digitization of over 1.5 million new observations from British Royal Naval ships logbooks from the 1914–1923 period, using volunteers to transcribe the hand-written pages. Projects such as Old Weather and the 20th Century Reanalysis (Compo *et al.* 2011) continue to refine our estimates and understanding of historical climate variability.

In the 75 years since Callendar (1938), great progress has been made in understanding the past changes in Earth's climate, and whether continued warming is beneficial or not. In 1938, Callendar himself concluded that, "*the combustion of fossil fuel [...] is likely to prove beneficial to mankind in several ways*", notably allowing cultivation at higher northern latitudes, and because, "*the return of the deadly glaciers should be delayed indefinitely*".

Callendar's achievements are remarkable, especially as he was an amateur climatologist, doing much of his research in his spare time, without access to a computer. He was meticulous in his approach, and a large collection of his notebooks are currently archived at the University of East Anglia. For completeness, Callendar's tabulated global land temperatures are presented as Supplementary Information.

Finally, Fleming (2007) gives a comprehensive review of the life and times of Guy Stewart Callendar (Fig. 2), whose efforts in attempting to estimate and understand the changing temperatures of the planet should be suitably acknowledged.

Acknowledgement

The authors would like to thank Colin Morice for providing the CRUTEM4 data, as well as John Kennedy, Vikki




Figure 2. A photograph of Guy Stewart Callendar (1898–1964), taken in 1934. Image from University of East Anglia archives. Additional photographs can be found in Fleming (2007).

Frith, Samantha Burgess, Keith Shine, Ian Strangeways, Steve Easterbrook, Geert Jan van Oldenborgh, and two anonymous reviewers for useful comments. EH is supported by a NERC Advanced Research Fellowship and PDJ has been supported by the US Dept of Energy (Grant DE-SC0005689).

References

- Arrhenius S. 1896. On the influence of carbonic acid in the air upon the temperature of the ground. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* **41**(251): 237–276.
- Bronnimann S. 2009. Early twentieth-century warming. *Nature Geoscience* **2**: 735–736, doi:10.1038/ngeo670.
- Callendar GS. 1938. The artificial production of carbon dioxide and its influence on temperature. *QJRMS* **64**(275): 223–240, doi:10.1002/qj.49706427503.
- Callendar GS. 1941. Infra-red absorption by carbon dioxide, with special reference to atmospheric radiation. *QJRMS* **67**: 263–275, doi:10.1002/qj.49706729105.
- Callendar GS. 1961. Temperature fluctuations and trends over the earth. *QJRMS* **87**(371): 1–12, doi: 10.1002/qj.49708737102.
- Compo GP, Whitaker JS, Sardeshmukh PD, Matsui N, Allan RJ, Yin X, Gleason BE, Vose RS, Rutledge G, Bessemoulin P, Brnnimann S, Brunet M, Crouthamel RI, Grant AN, Groisman PY, Jones PD, Kruk MC, Kruger AC, Marshall GJ, Mauerer M, Mok HY, Nordli, Ross TF, Trigo RM, Wang XL, Woodruff SD, Worley SJ. 2011. The twentieth century reanalysis project. *Quarterly Journal of the Royal Meteorological Society* **137**(654): 1–28, doi:10.1002/qj.776.

- Delworth TL, Knutson TR. 2000. Simulation of early 20th century global warming. *Science* **287**: 2246–2250, doi:10.1126/science.287.5461.2246.
- Ekholm N. 1901. On the variations of the climate of the geological and historical past and their causes. *QJRMS* **27**: 1–61.
- Fleming J. 2007. *The Callendar effect: the life and times of Guy Stewart Callendar (1898-1964), the scientist who established the carbon dioxide theory of climate change*. American Meteorological Society.
- Fourier J. 1824. Rapport sur la temperature du globe terrestre et sur les spaces planétaires. *Mémoires Acad. Royale des Sciences de L'Institut de France* **1824**: 590–604.
- Jones PD, Lister DH, Osborn TJ, Harpham C, Salmon M, Morice CP. 2012. Hemispheric and large-scale land-surface air temperature variations: An extensive revision and an update to 2010. *Journal of Geophysical Research* **117**: D05 127, doi:10.1029/2011JD017139.
- Jones PD, Osborn TJ, Briffa KR. 1997. Estimating sampling errors in large-scale temperature averages. *J. Climate* **10**: 2548–2568.
- Jones PD, Wigley TML. 2010. Estimation of global temperature trends: What's important and what isn't. *Climatic Change* **100**: 59–69, doi:10.1007/s10584-010-9836-3.
- Köppen W. 1881. Über mehrjährige perioden der witterung—III. Mehrjährige änderungen der temperatur 1841 bis 1875 in den tropen der nördlichen und südlichen gemässigten zone, an den Jahresmitteln. untersucht. *Zeitschrift der Österreichischen Gesellschaft für Meteorologie* : 141–150.
- Le Treut H, Somerville R, Cubasch U, Ding Y, Mauritzen C, Mokssit A, Peterson T, Prather M. 2007. *Historical overview of climate change*. In: *Climate Change 2007: The physical science basis*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Madden RA, Shea DJ, Branstator GW, Tribbia JJ, Weber R. 1993. The effects of imperfect spatial and temporal sampling on estimates of the global mean temperature: Experiments with model and satellite data. *J. Climate* **6**: 1057–1066.
- Parker DE. 2010. Urban heat island effects on estimates of observed climate change. *WIREs: Climate Change* **1**(1): 123–133, doi:10.1002/wcc.21.
- Plass GN. 1956. The carbon dioxide theory of climatic change. *Tellus* **8**: 140–154.
- Tyndall J. 1861. On the absorption and radiation of heat by gases and vapours, and on the physical connexion of radiation, absorption, and conduction. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* **22**(146): 169–194.
- Weart S. 2008. *The discovery of global warming*. Harvard University Press, USA.
- Wood KR, Overland JE, Jónsson T, Smoliak BV. 2010. Air temperature variations on the Atlantic-Arctic boundary since 1802. *Geophysical Research Letters* **37**: L17 708, doi:10.1029/2010GL044176.